

PUPT-1389, IASSNS-HEP-93/16

March 1993

Emergence of Coherent Long Wavelength Oscillations After a Quench: Application to QCD

KRISHNA RAJAGOPAL[★]

*Department of Physics
Joseph Henry Laboratories
Princeton University
Princeton, N.J. 08544*

FRANK WILCZEK[†]

*School of Natural Sciences
Institute for Advanced Study
Olden Lane
Princeton, N.J. 08540*

★ Research supported in part by a Natural Sciences and Engineering Research Council of Canada 1967 Fellowship and in part by a Charlotte Elizabeth Procter Fellowship. RAJAGOPAL@PUPGG.PRINCETON.EDU

† Research supported in part by DOE grant DE-FG02-90ER40542. WILCZEK@IASSNS.BITNET

ABSTRACT

To model the dynamics of the chiral order parameter in a far from equilibrium phase transition, we consider quenching in the $O(4)$ linear sigma model. We argue, and present numerical evidence, that in the period immediately following the quench long wavelength modes of the pion field are amplified. This results in large regions of coherent pion oscillations, and could lead to dramatic phenomenological consequences in heavy ion collisions.

Among the most interesting speculations regarding ultra-high energy hadronic or heavy nucleus collisions is the idea that regions of misaligned vacuum might occur [1,2,3,4,5]. In such misaligned regions, which are analogous to misaligned domains in a ferromagnet, the chiral condensate points in a different direction from that favored in the ground state. If we parametrize the condensate using the usual variables of the sigma model, misaligned vacuum regions are places where the four-component $(\sigma, \vec{\pi})$ field, that in the ground state takes the value $(v, 0)$, is partially aligned in the π directions. If they were produced, misaligned vacuum regions plausibly would behave as “pion lasers”, relaxing to the ground state by coherent pion emission. They would produce clusters of pions bunched in rapidity with highly non-Gaussian charge distributions. Thus a misaligned vacuum region starting with the field in the $\pi_1 - \pi_2$ plane would emit only charged pions (equally positive and negative, since the fields are real), while a misaligned vacuum region starting with the field pointing in the π_3 direction would emit only neutral pions. More generally if we define

$$R \equiv \frac{n_{\pi^0}}{n_{\pi^0} + n_{\pi^+ \pi^-}} \quad (1)$$

then assuming that all initial values on the 3-sphere are equally probable, the probability distribution $\mathcal{P}(R)$ is given by

$$\mathcal{P}(R) = \frac{1}{2} R^{-1/2} . \quad (2)$$

As an example of (2), we note that the probability that the neutral pion fraction R is less than .01 is 0.1! This is a graphic illustration of how different (2) is from what one would expect if individual pions were independently emitted with no isospin correlations. According to (2) there is a substantial probability that, say, a cluster of 70 pions would all be charged – something that could essentially never occur for incoherent emission. There may be some hint of such behavior in the Centauro (overwhelmingly charged) and anti-Centauro (overwhelmingly neutral) events reported in the cosmic ray literature.[6,7] Here we propose a concrete

mechanism by which such phenomena might arise in heavy ion collisions, for which the plasma is far from thermal equilibrium.

In previous work [5,8] we considered the equilibrium phase structure of QCD. We argued that QCD with two massless quark flavors probably undergoes a second-order chiral phase transition. For many purposes it is a good approximation to treat the u and d quarks as approximately massless, and we expect that real QCD has a smooth but perhaps rapid transition as a function of decreasing temperature from small intrinsic to large spontaneous chiral symmetry breaking. At first sight it might appear that a second-order phase transition is especially favorable for the development of large regions of misaligned vacuum. Indeed the long-lived, long-wavelength critical fluctuations which provide the classic signature of a second-order transition *are* such regions. Unfortunately the effect of the light quark masses, even though they are formally much smaller than intrinsic QCD scales, spoil this possibility [5]. The pion masses, or more precisely the inverse correlation length in the pion channel, are almost certainly *not* small compared to the transition temperature. As a result the misaligned regions are modest affairs at best even near the critical temperature. They extend at most over a few thermal wavelengths, and almost certainly do not contain sufficient energy to radiate large numbers of correlated pions.

Here we will consider an idealization that is in some ways opposite to that of thermal equilibrium, that is the occurrence of a sudden quench from high to low temperatures, in which the $(\sigma, \vec{\pi})$ fields are suddenly removed from contact with a high temperature heat bath and subsequently evolve mechanically. We shall show that long wavelength fluctuations of the light fields (the pions in QCD, which would be massless if not for the quark masses) can develop following a quench from some temperature $T > T_c$ to $T = 0$. The long wavelength modes of the pion fields are unstable and grow relative to the short wavelength modes.

Before we enter into the details of our model and simulations it seems appropriate to discuss the qualitative reason for the result just mentioned, which we suspect

may be of wider interest. Whenever one has spontaneous breaking of a continuous global symmetry, massless Nambu-Goldstone bosons occur. The masslessness of these modes occurs through a cancellation following the schema

$$m^2 = -\mu^2 + \lambda\phi^2, \quad (3)$$

where the second term arises from interaction with a condensate whose expectation value $\langle\phi\rangle^2$ satisfies $\langle\phi\rangle^2 = \mu^2/\lambda \equiv v^2$ in the ground state. However following a quench the condensate starts with its average at 0 rather than v , and generally oscillates before settling to its final value. Whenever $\langle\phi\rangle^2 < v^2$, m^2 will be negative, and *sufficiently long wavelengths fluctuations in the Nambu-Goldstone boson field will grow exponentially*. The same mechanism will work, though less efficiently, if one has only an approximate symmetry and approximate Nambu-Goldstone modes, or for that matter other modes whose effective m^2 is ‘accidentally’ pumped negative by their interactions with the condensate.

Returning to QCD, we shall use the classic Gell-Mann–Levy Lagrangian [9] to describe the low energy interactions of pions:

$$\mathcal{L} = \int d^4x \left\{ \frac{1}{2} \partial^i \phi^\alpha \partial_i \phi_\alpha - \frac{\lambda}{4} (\phi^\alpha \phi_\alpha - v^2)^2 + H\sigma \right\}, \quad (4)$$

where ϕ is a four-component vector in internal space with entries (σ, π) . Here, λ , v , and $H \propto m_q$ are to be thought of as parameters in the low energy effective theory obtained after integrating out heavy degrees of freedom. We shall treat (4) as it stands as a classical field theory, since the phenomenon we are attempting to address is basically classical and because as a practical matter it would be prohibitively difficult to do better. We shall be dealing with temperatures of order 200 MeV or less, so that neglect of heavier fields seems reasonable.

To model a quench, we begin at a temperature well above T_c . The typical configurations have short correlation lengths and $\langle\phi\rangle \sim 0$. ($\langle\phi\rangle \neq 0$ because $H \neq 0$.) One then takes the temperature instantaneously to zero. The equilibrium configuration is an ordered state with the ϕ field aligned in the σ direction throughout

space, but this is not the configuration in which the system finds itself. The actual, disordered configuration then evolves according to the zero temperature equations of motion obtained by varying (4). Quenching in magnet models has been much studied in condensed matter physics [10]. However in that context it is usually appropriate to use diffusive equations of motion, because the magnet is always in significant contact with other light modes (*e.g.* phonons). For this reason the condensed matter literature we are aware of does not directly apply to our problem.

Let us now consider how the physics of quenching may be applied in relativistic heavy ion collisions. In Bjorken's [11] picture of such a collision, the incident nuclei as seen in the center of mass frame are both Lorentz contracted into pancake shapes. They pass through each other, and leave behind a region of hot vacuum. The baryon number of the incident nuclei ends up in that part of the plasma heading approximately down the beam pipe, and the central rapidity region which we consider in this paper consists of plasma with approximately zero baryon number which expands and cools through $T = T_c$ and eventually hadronizes into the detected pions. If the collision is energetic enough to create a region of plasma well above $T = T_c$, the ϕ field will indeed be fluctuating among an ensemble of disordered configurations. The plasma cools rapidly as it expands, and cannot be exactly in thermal equilibrium. If it cools fast enough, the configuration of the ϕ field will "lag," and as in a quench the system will find itself more disordered than the equilibrium configuration appropriate to the current temperature.

We have done numerical simulations of quenching to zero temperature in the linear sigma model with an explicit symmetry breaking term $H\sigma$ which makes the pions massive. Turok and Spergel [12] have considered this scenario with no explicit symmetry breaking term as a cosmological model for large scale structure formation in the early universe. They find a scaling solution in which the size of correlated domains grows at the speed of light as larger and larger regions come into causal contact and align. When the $O(N)$ symmetry is explicitly broken, however, we do not expect a scaling solution. The $H\sigma$ term tilts the potential, the vacuum manifold is not degenerate, and in a time of order m_π^{-1} the scalar field in

all regions (whether in causal contact or not) will be oscillating about the sigma direction.

In numerically simulating a quench, we choose ϕ and $\dot{\phi}$ randomly independently on each site of a cubic lattice. This means that the lattice spacing a represents the correlation length in the initial conditions. If the initial conditions are chosen to model a thermal ensemble at some temperature $T > T_c$, then the $O(N)$ symmetry is not spontaneously broken, and the lattice spacing a represents the π and σ correlation lengths which are approximately degenerate. Because the initial conditions must not distinguish between the σ and π directions in internal space, we must use the linear sigma model (4) instead of integrating out the σ which is heavy at zero temperature to obtain the nonlinear sigma model. In the simulation whose results are shown in Figure 1, we chose ϕ and $\dot{\phi}$ randomly from gaussian distributions centered around $\phi = \dot{\phi} = 0$ and with root mean square variance $v/2$ and v respectively. The three parameters v , H , and λ in (4) determine m_π , m_σ , and $f_\pi = \langle 0|\sigma|0\rangle$ according to

$$\lambda \langle 0|\sigma|0\rangle \left(\langle 0|\sigma|0\rangle^2 - v^2 \right) - H = 0 \quad , \quad (5)$$

$$m_\pi^2 = \frac{H}{\langle 0|\sigma|0\rangle} \quad , \quad \text{and} \quad m_\sigma^2 = 3\lambda \langle 0|\sigma|0\rangle^2 - \lambda v^2 \quad . \quad (6)$$

Note that $f_\pi = \langle 0|\sigma|0\rangle > v$ for $H \neq 0$. In interpreting our results we must remember that while in the code we are free to choose the energy scale by setting the lattice spacing $a = 1$, a actually represents the initial correlation length. In choosing the parameters for Figure 1 we assumed that $a = (200 \text{ MeV})^{-1}$ and then chose $v = 87.4 \text{ MeV} = 0.4372 \text{ } a^{-1}$, $H = (119 \text{ MeV})^3 = 0.2107 \text{ } a^{-3}$, and $\lambda = 20.0$ so that $f_\pi = 92.5 \text{ MeV}$, $m_\pi = 135 \text{ MeV}$, and $m_\sigma = 600 \text{ MeV}$.

With parameters and initial conditions chosen, we evolve the initial configuration according to the equations of motion using a standard finite difference, staggered leapfrog scheme. (For details of the numerical method see [13,14] and in particular equations 32-35 of [14].) We used a time step $dt = \min(\frac{a}{10}, \frac{2\pi}{10m_\sigma})$. We

verified that our results do not change if the time step is reduced. After each two time steps, we computed the spatial fourier transform of the configuration, and from that obtained the angular averaged power spectrum. In Figure 1, we plot the power in modes of the pion and sigma fields with spatial wave vectors of several different magnitudes k as a function of time. All the curves start at approximately the same value at $t = 0$ because the initial power spectrum is white since we chose ϕ independently at each lattice site. The behaviour of the low momentum pion modes is striking. While the initial power spectrum is white and, as ergodicity arguments would predict, the system at late times is approaching an equilibrium configuration in which the equipartition theorem holds, at intermediate times of order several times m_π^{-1} the low momentum pion modes are oscillating coherently in phase with large amplitudes.

The simulation was done in a 64^3 box. We verified that finite size effects are not important even for the longest wavelength ($ka = 0.20$) mode shown in Figure 1 by checking that the behaviour of the $ka = 0.31$ mode is the same in a 32^3 and a 64^3 box. The behaviour of only one component of the pion field is shown in Figure 1a. The other two look qualitatively the same. It should be noted that the exact height of the peaks in the curves depend on the specific initial conditions. If the simulation is run with the same initial distribution for ϕ and $\dot{\phi}$ but with a different seed for the random number generator, the heights of the peaks change, and the relative sizes of the peaks in the three different pion directions change. The qualitative features of Figure 1 — the growth of long wavelength modes of the pion field — do not depend on the specific realization of the initial conditions. We discuss below how the results change as a function of parameters in the potential and the initial distributions.

Let us compare the growth of long wavelength modes found numerically with expectations from the mechanism previously discussed. Suppose the potential V in (4) were simply $V(\phi) = (m^2/2)\phi^\alpha\phi_\alpha$. Then, the equations of motion would be linear, and modes with different spatial wave vector $\vec{k}$ would be uncoupled. Each curve in Figure 1 would be a sinusoid with period $\pi/\sqrt{m^2 + \vec{k}^2}$ and constant

amplitude. (The power spectrum, being quadratic in the fields, oscillates with one half the period of the fields.) The period of the oscillations in Figure 1a is indeed given by $\pi/\sqrt{m_\pi^2 + \vec{k}^2}$, but the amplitudes are far from constant. This behaviour can be qualitatively understood by approximating ϕ^2 in the nonlinear term in the equation of motion by its spatial average:

$$\phi^\alpha \phi_\alpha(\vec{x}, t) \sim \langle \phi_\alpha \phi^\alpha \rangle(t) . \quad (7)$$

This approximation is exact in the large N limit [12]. In our problem, $N = 4$. Using (7) and doing the spatial fourier transform, the equation of motion for the pion field becomes

$$\frac{d^2}{dt^2} \vec{\pi}(\vec{k}, t) = -m_{eff}^2(k, t) \vec{\pi}(\vec{k}, t) \quad (8)$$

with the time dependent “mass” given by

$$m_{eff}^2(k, t) \equiv -\lambda v^2 + k^2 + \lambda \langle \phi^2 \rangle(t) \quad (9)$$

where $k = |\vec{k}|$.

Figure 2 shows the time evolution of $\langle \phi^2 \rangle$ in the same simulation whose results are shown in Figure 1. In the initial conditions, ϕ is gaussian distributed with $\langle \phi^2 \rangle < v^2$. Therefore, for a range of wavevectors with k less than some critical value, $m_{eff}^2 < 0$ and the long wavelength modes of the pion field start growing exponentially. $\langle \phi^2 \rangle$ grows and then executes damped oscillations about its ground state value $\langle 0|\sigma|0 \rangle^2$. A wave vector k mode of the pion field is unstable and grows exponentially whenever $\langle \phi^2 \rangle < v^2 - k^2/\lambda$. Modes with $k^2 > \lambda v^2$ can never be unstable. The $k = 0$ mode is unstable during the periods of time when the $\langle \phi^2 \rangle$ curve in Figure 2 is below v^2 . Since $\langle \phi^2 \rangle$ is oscillating about $\langle 0|\sigma|0 \rangle^2 > v^2$, after some time the oscillations have damped enough that $\langle \phi^2 \rangle$ never drops below v^2 , and from that time on all modes are always stable and oscillatory. In general, longer wavelength modes are unstable for more and for longer intervals of time

than shorter wavelength modes as $\langle\phi^2\rangle$ oscillates. Also, m_{eff}^2 is more negative and more growth occurs for modes with smaller k . Making the approximation (7) and thus using (8) cannot be expected to completely reproduce the effects of the nonlinear term in (4) which is local in position space. Nevertheless, it predicts that the long wavelength modes of the pion field go through alternating periods of oscillatory behaviour and exponential growth spurts and therefore gives us a good understanding of the behaviour of these modes in Figure 1a. Because the timing of the growth spurts for different (low) k modes are all determined by $\langle\phi^2\rangle$, different modes have their growth spurts at the same times and all the long wavelength modes in Figure 1a oscillate in phase. The longer wavelength modes have more, stronger, and longer growth spurts, and are therefore amplified more as in Figure 1a. In an equilibrium phase transition, explicit symmetry breaking keeps the correlation length at T_c finite and, in the case of QCD, too short to be of interest.[5] After a quench, on the other hand, arbitrarily long wavelength modes of the pion field are amplified even though the pion mass is non-zero.

What about the σ modes? For $\sigma^2 < v^2/3$, the effective mass² for the $k = 0$ mode is negative. This condition is far less likely to be satisfied than $\langle\phi^2\rangle < v^2$. Hence, while some growth of the low momentum σ modes is possible, as Figure 1 confirms, they do not grow as much as the low momentum pion modes do. The behaviour of the sigma modes is also complicated by the fact that oscillations with two periods — $2\pi/m_\pi$ and $2\pi/m_\sigma$ — contribute. Roughly, this occurs because oscillations ostensibly along the pion directions are oscillations in a curved valley and are therefore seen in sigma also.

At late times $\langle\phi^2\rangle$ oscillates with small enough amplitude that it never goes below v^2 and consequently no modes are ever unstable. Hence, if we make the approximation (7) we would expect that at late times each mode in Figure 1a would continue to oscillate with approximately constant amplitude, with the longer wavelength modes maintaining the large amplitudes acquired during their exponential growth spurts. That this is not what is seen in Figure 1a reflects the effects we neglected in making the approximation (7). Because the modes are in fact cou-

pled and the equations of motion are actually nonlinear, ergodicity arguments suggest that eventually equipartition should apply. If the energy is equally divided among modes, then the (amplitude)² in a mode should be inversely proportional to $(m_\pi^2 + k^2)$. Also, the amplitude of the sigma modes at late times should be less than that of pion modes of the same k because $m_\sigma > m_\pi$. The results shown in Figure 1 are consistent with the assumption that at late times equipartition applies, although at $t = 80a$ the longest wavelength pion modes are still decreasing in amplitude. (We have verified that at even later times they do reach equipartition.) It is reasonable that longer wavelength modes take longer to decrease in amplitude than shorter wavelength modes. As the system heads toward equipartition, power has to flow from low momentum modes to higher momentum modes. This plausibly happens more quickly at higher momentum because higher momentum modes are coupled to a larger number of modes, or, in the continuum, a larger volume of momentum space. Hence, we end up with the striking phenomena of Figure 1a: while the system begins and ends with small $\vec{\pi}(\vec{k}, t)$, in between the long wavelength pion modes are dramatically amplified.

In modelling a heavy ion collision, we should include the effects of the expansion of the plasma after a quench. In an equilibrium phase transition, the expansion causes a decrease in the temperature. After a quench, however, $T = 0$ and the expansion has the effect of reducing the energy per unit volume in the system. The simplest way to model this is to introduce a time dependent lattice spacing $a(t)$. This introduces a $\dot{\phi}a/a$ term in the equations of motion which, for $a(t)$ increasing with time, leads to damping. Doing this, we found that each of the modes (now defined by spatially fourier transforming from comoving $\vec{x}$ to comoving $\vec{k}$) red shifts with time. However, as long as the $\dot{\phi}$ term does not dominate over the $\ddot{\phi}$ term low momentum modes still become unstable and grow with respect to high momentum modes. In a real heavy ion collision, the expansion is anisotropic. The plasma expands much more rapidly along the beam direction than in the transverse directions [11]. While we have not attempted to model this explicitly, it probably means that correlated volumes do not grow large in the longitudinal

direction. The large expansion rate will rapidly damp modes of the pion field with wave vectors in the longitudinal direction. However, it seems possible that for a particular longitudinal position (or, equivalently [11], a particular rapidity) coherent fluctuations on length scales all the way up to the transverse extent of the plasma could develop. Another effect of the expansion is that after some time (perhaps of order $10 \text{ fm}^{-1} \sim 7m_\pi^{-1}$ [11]) the energy density is low enough that the description in terms of classical fields no longer makes sense. After this time, one has individual pions flying off towards the detector. From Figure 1a, it is plausible that at this time there are large long wavelength oscillations of the pion field.

If the initial conditions or parameters are such that the initial value of the mean energy per unit volume E is much more than $\frac{\lambda}{4}v^4$ no striking growth of low momentum modes occurs. Indeed when the total energy is large, the field will be able to climb far up the potential to $|\phi| \gg v$, and therefore $\langle \phi^2 \rangle(t)$ will always be greater than v^2 , and no modes will become unstable. Requiring the energy not to be too large compared to the height of the potential at $\phi = 0$ can be seen either as a constraint on the parameters of the potential ($V(\phi = 0) \sim \frac{\lambda}{4}v^4$) for given initial conditions or on the initial conditions for a given potential. In the simulation of Figure 1, the total energy $E \sim 2\frac{\lambda}{4}v^4$. As parameters (or initial conditions) are changed so that the total energy is increased, the growth of low momentum modes becomes less and less prominent, and when $E \gtrsim 10\frac{\lambda}{4}v^4$, dramatic growth does not occur.

As long as ϕ is centered around $\phi = 0$ initially, and as long as E is not too large, the choice of initial distributions has little qualitative effect. We have tried distributions ranging from $\phi \equiv 0$ with $\dot{\phi}$ chosen from a gaussian distribution to $\dot{\phi} \equiv 0$ with ϕ chosen from a uniform distribution with $\phi^2 \leq v^2$, and in all cases obtained results similar to those of Figure 1.

Let us attempt to estimate the conditions that might govern a realistic quench. We will make a crude estimate of the lattice spacing a and initial distributions for ϕ and $\dot{\phi}$ using the one loop temperature dependent effective potential with $H = 0$.

At large temperatures, this is given by [15,16]

$$V(\phi, T) = m^2(T)\phi^2 + \frac{\lambda}{4}\phi^4 \quad (10)$$

where the temperature dependent mass is given by

$$m^2(T) = \lambda(-v^2 + T^2/4) \quad (11)$$

While this is a crude approximation, it gives $T_c = 2v = 175$ MeV which is coincidentally quite close to the lattice gauge theory value [17]. If we quench from an initial temperature T_i , the lattice spacing will be given by the correlation length at T_i

$$a = m^{-1}(T_i) \quad (12)$$

For $T_i = 1.2 T_c = 2.4 v$, this gives $a^{-1} = 2.97 v = 260$ MeV.

We will model initial conditions with correlation length a by choosing ϕ (and $\dot{\phi}$) independently at each lattice site. ϕ should therefore be distributed according to the probability distribution

$$\exp - \frac{a^3}{T_i} \left(\frac{1}{2} m^2(T_i) \phi^2 + \frac{\lambda}{4} \phi^4 \right) \quad (13)$$

Because it is cut off by the quartic term, this distribution has much less weight at large $|\phi|$ than a gaussian. We approximate it by the truncated gaussian distribution

$$\exp - \left(\frac{1}{2} m^2(T_i) \phi^2 \right) \text{ for } |\phi| < \phi_{max} ; \quad 0 \text{ for } |\phi| > \phi_{max} \quad (14)$$

where ϕ_{max} is chosen so that ϕ^2 has the same expectation value in the distributions (13) and (14). To choose initial conditions for $\dot{\phi}$, we note that $\langle \dot{\phi}^2 \rangle = \langle \phi^2 \rangle$ in lattice units. (This is true for a quadratic potential, and is also true in general if equipartition applies.) Hence, we use the same distribution for $\dot{\phi}$ as for ϕ . We now have a crude recipe for choosing initial conditions appropriate to model a quench from an initial temperature T_i to $T = 0$.

What does our crude recipe predict? From (13), we expect that in lattice units $\phi_{max} \sim (\lambda/2)^{1/4} = 0.56$. In fact, for $T_i = 1.2 T_c$, we obtain $\phi_{max} = 0.63 a^{-1} = 1.87 v$. The value of ϕ_{max} is crucial, because it determines over how much of the potential ϕ is distributed. We see that choosing ϕ gaussian distributed with variance $v/2$ as we did in Figure 1 was overly optimistic. For $T_i = 1.2 T_c$, it is more reasonable to assume that ϕ is distributed according to (14) with $\phi_{max} = 1.87 v$. With these initial conditions, the energy per unit volume turns out to be $E \sim 8 \frac{\lambda}{4} v^2$ and the pion field behaves as shown in Figure 3. Low momentum modes do indeed grow, but less dramatically than in Figure 1. Also, the system finds its way to a final state described by equipartition more quickly than in Figure 1. Nevertheless, there is a period of time after the quench when low momentum modes have much more power than higher momentum modes.

If T_i is taken to be higher, ϕ_{max}/v and $1/av$ increase, $E/(\frac{\lambda}{4}v^4)$ increases, and the phenomenon of interest is washed out. It is not unreasonable, however, to consider quenching from temperatures larger than but comparable to T_c . If the initial temperature were much higher, the system could plausibly stay in thermal equilibrium until close to T_c . Hence, choosing $T_i = 1.2 T_c$ and thus obtaining parameters and initial distributions like those of Figure 3 may be a reasonable if crude approximation to a real heavy ion collision.

We have made many idealizations and approximations. Nevertheless, it seems possible to us that the essential qualitative feature of the phenomenon we have elucidated — long wavelength pion modes experiencing periods of negative mass² and consequent growth following a quench — could occur in real heavy ion collisions. Given the explicit symmetry breaking, one might have expected the dynamics following a quench from a “generic” initial state to be featureless. The mechanism here discussed provides a robust counterexample that should be applicable in other contexts besides the QCD phase transition. Examples could include the reheating of some inflationary universe models and the quenching of spin systems which order at low temperatures, for which the interactions of the spins with phonons are less important than their interactions among themselves. If a heavy ion collision

is energetic enough that there is a central rapidity region of high energy density and low baryon number, and if such a region cools rapidly enough that it can be modelled as a quench, this will be detected by observing clusters of pions of similar rapidity in which the the fraction of neutral pions is fixed. This ratio will be different in different clusters and will follow a distribution like (2). We can think of no process besides a QCD phase transition with the chiral order parameter far out of equilibrium that could produce such a signature.

Acknowledgements: We are grateful to David Spergel and Neil Turok for giving us the code created by the authors of [12, 13, and 14], and to both of them and Ue-Li Pen for many very helpful conversations.

REFERENCES

1. A. Anselm and M. Ryskin, *Phys. Letters* **B226**, 482 (1991).
2. J.-P. Blaizot and A. Krzywicki, *Phys. Rev.* **D46**, 246 (1992).
3. J. D Bjorken, *Int. J. Mod. Phys.* **A7**, 4189 (1992); J. D. Bjorken, *Acta Physica Polonica* **B23**, 561 (1992).
4. K. L. Kowalski and C. C. Taylor, *Disoriented Chiral Condensate: A White Paper for the Full Acceptance Detector*, CWRUTH-92-6, hep-ph/9211282, 1992.
5. K. Rajagopal and F. Wilczek, *Static and Dynamic Critical Phenomena at a Second Order QCD Phase Transition* PUPT-1347, IASSNS-HEP-92/60, hep-ph/9210253, to appear in *Nucl. Phys.* **B**.
6. C. M. G. Lattes, Y. Fujimoto, and S. Hasegawa, *Phys. Rept.* **65**, 151 (1980).
7. J. J. Lord and J. Iwai, Univ. of Washington preprint (paper 515 submitted to the International Conference on High Energy Physics, Dallas, August 1992); J. Iwai (JACEE collaboration), UWSEA 92-06.
8. F. Wilczek, *Int. J. Mod. Phys.* **A7**, 3911 (1992). This elaborates earlier work of R. Pisarski and F. Wilczek, *Phys. Rev.* **D29**, 338 (1984).
9. M. Gell-Mann and M. Levy, *Nuovo Cimento* **16**, 705 (1960).
10. A. J. Bray, *Phys. Rev.* **B41**, 6724 (1990); T. J. Newman, A. J. Bray, and M. A. Moore, *Phys. Rev.* **B42**, 4514 (1990); and references therein.
11. J. D. Bjorken, *Phys. Rev.* **D27**, 140 (1983).
12. N. Turok and D. N. Spergel, *Phys. Rev. Lett.* **66**, 3093 (1991).
13. D. N. Spergel, N. Turok, W. H. Press, and B. S. Ryden, *Phys. Rev.* **D43**, 1038 (1991).
14. W. H. Press, B. Ryden, and D. N. Spergel, *Astrophys. J.* **347**, 590 (1989).
15. L. Dolan and R. Jackiw, *Phys. Rev.* **D9**, 3320 (1974).

16. C. Contreras and M. Loewe, *Int. J. Mod. Phys.* **A5**, 2297 (1990).
17. C. Bernard *et al.*, *Phys. Rev.* **D45**, 3854 (1992).

FIGURE CAPTIONS

- 1) Time evolution of the power spectrum of the pion field and the sigma field at several spatial wavelengths after a quench at $t = 0$. ϕ and $\dot{\phi}$ were chosen at time $t = 0$ independently at each lattice site from gaussian distributions centered at the origin as described in the text. The parameters of the potential were chosen so that for a lattice spacing $a = (200 \text{ MeV})^{-1}$, the physical values $m_\pi = 135 \text{ MeV}$, $m_\sigma = 600 \text{ MeV}$, and $f_\pi = 92.5 \text{ MeV}$ are obtained. The simulation was performed on a 64^3 lattice and the time step was $a/10$. After every two time steps, the spatial fourier transform of ϕ was computed. For each component of ϕ , the power in all the modes with $k \equiv |\vec{k}|$ in bins of width $0.057a^{-1}$ were averaged. Figure 1a shows the time evolution of one component of the pion field. The curves plotted are (top to bottom in the figure) the average power in the modes in the momentum bins centered at $ka = 0.20, 0.26, 0.31, 0.37, 0.48, 0.60, 0.71, 0.94, 1.16, 1.39$, and 1.84 . The initial power spectrum is white and all the curves start at $t = 0$ at approximately 0.01 in lattice units. Hence, the longest wavelength pion modes are amplified by a factor of order 1000 relative to the shortest wavelength modes which are not amplified at all. Figure 1b shows the time evolution of the sigma field. Only four modes are plotted — $ka = 0.20, 0.48, 0.94$, and 1.84 . The vertical scale is different than in Figure 1a — the sigma modes do not grow nearly as much as the pion modes.
- 2) Time evolution of the spatially averaged $\langle \phi^2 \rangle$ for the same simulation whose results are shown in Figure 1. The horizontal line is at $\langle \phi^2 \rangle = v^2$.
- 3) Same as for Figure 1a, except that in this figure we have used $a^{-1} = 260 \text{ MeV}$ and have chosen initial conditions from the distribution (14) following the crude recipe outlined in the text to model a quench from $T_i = 1.2 T_c$ to $T = 0$. Note that the vertical scale is different than in Figure 1a. All the curves start at about 0.03 at $t = 0$. With these more realistic initial conditions, long wavelength modes of the pion field grow less than in Figure 1a, but are

still significantly amplified relative to shorter wavelength modes.

FIGURE 1a




FIGURE 1b




FIGURE 2




FIGURE 3

